

Toxicological Studies and Biodegradability Prediction of Isonicotinylhydrazide

Kamel Mokhnache*, Soraya Madoui and Nouredine Charef

Laboratory of Applied Biochemistry, University Ferhat Abbas Setif 1,19000, Algeria.

*Corresponding Author E-mail:kamelmokhnache@yahoo.com

Received: December 28, 2018; **Accepted:** January 6, 2019; **Published:** January 12, 2019

Abstract: The antitubercular drug isonicotinylhydrazide (Isoniazid) was selected for toxicological and biodegradation studies. In silico ecotoxicological and biodegradability predictions were carried out online computer software programs such as Pro Tox, Pred-skin, Endocrine Disruptor Knowledge Base (EDKB) and UM-BBD. In silico predictions, results showed high toxicity of this drug; hepatotoxicity with probability of 93%, carcinogenicity; 81% and without nephrotoxic effect. For biodegradability prediction, the results indicate the ability of microorganism to degrade the drug with no-toxic resultant products. We conclude the possibility to using the drug with high precautions from environment and human health.

Keywords: toxicological, biodegradability, microorganisms, online software programs.

Citation: Kamel Mokhnache, Soraya Madoui and Nouredine Charef. 2019. Toxicological Studies and Biodegradability Prediction of Isonicotinylhydrazide. International Journal of Current Innovations in Advanced Research, 2(1): 1-5.

Copyright: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. **Copyright©2019;** Kamel Mokhnache, Soraya Madoui and Nouredine Charef.

Methods

Online software programs

ProTox web server was used for toxicity prediction method. The University of Minnesota Biocatalysis/Biodegradation Database (UM-BBD, <http://umbbd.msi.umn.edu/>) was used for predict microbial biocatalytic reactions and biodegradation pathways.

Results and Discussion

Toxicological studies

Results displayed in Table 1, showed high toxicity of Isoniazid; hepatotoxicity with probability of 93%, carcinogenicity; 81% and without nephrotoxic effect.

Table 1. Toicity evaluation of isoniazid

Toxicity	Prediction	Probability %
Hepatotoxicity	Active	93
Carcinogenicity	Active	81
Nephrotoxicity	Inactive	-

Bioderadability prediction

Numerous chemicals used in human-life are supposed to be pollutants [1], when were accumulated in environment, interacted with biological system caused mortal interactions.

Thus, obliged to create processes for decrease these pollutants and guard living organism. Among of these processes; the biodegradability [2], the ability of soil bacteria to decompose organic matter by extracellular enzymes production, led to facilitate and accelerate the mineralization process [3].

Several studies, confirmed the accompaniment of both increasing of microorganisms and the rate of mineralization when the organic matter introduced in the soil [4]. Recently, the progress in computational studies led to developed different rules of biotransformation of organic groups from UM-BBD data.

Results displayed in Figure 1; represents various pathways of isoniazid biodegradation and the resultant products. We can summarize isoniazid biodegradation in the following steps by different pathways:

In first time, isoniazid was biodegraded to five compounds (2 to 6), with oxidative and reductive reactions of bacterial enzymes such as monooxygenases, dioxygenases, hydrolases and nitroreductases.

bt0042: From Nitrobenzene to Catechol [5] by dioxygenase (2), bt0067: From secondary Amide to Carboxylate + primary Amine [6] by hydrolase (3, 4, 7, 13,17, 24, 25,35), bt0078: From Nitrobenzenoid to Hydroxylaminobenzenoid [8] by nitroreductase (5), bt0080: From Nitroaromatic to Aminoaromatic [9] by NAD (P) H nitroreductase (6).

After these steps, each one of these compounds was transformed to several compounds until obtain degradation final compounds with the following reactios pathways:

bt0037: From Hydroxylaminobenzene to 2-Aminophenol [10] by hydroxylaminobenzene mutase (15, 16), bt0035: from aromatic Hydroxylamine to aromatic Amine [10] by reductase (14), bt0042: from 1-Nitro-2-unsubstituted aromatic to *vic*-Dihydroxyaromatic [10] by nitrobenzene reductase (7), bt0051: from isonicotinate to pyridine [11] by isonicotinate decarboxylase (12), bt0005: from *vic*-unsubstituted Aromatic to *vic*-Dihydroxyaromatic [12] by dioxygenase (11), bt0065: from 1-Amino-2-unsubstituted aromatic to *vic*-Dihydroxyaromatic [13] by dioxygenase (7), bt0078: from Nitrobenzenoid to Hydroxylaminobenzenoid [14] by nitroreductase (13), bt0254: from *vic*-Dihydroxyaromatic to intradiol ring cleavage [15] by dioxygenase (8,17). bt0351: from *vic*-Dihydroxybenzenoid to 2-Oxopent-4-enoate derivative + Carboxylate [16] (9, 10, 24), bt348: from [2-halo]Maleylacetate derivative to Succinate (18,19, 27), bt0377: from 3-Hydroxy-2-methylpyridine-5-carboxylate derivative to 2-(Acetamidomethylene)succinate derivative [17] by oxygenase, bt0318: from Carbamate to Amine [18] (31), bt0373 from From 3,4-Dehydroadipyl-CoA to Succinyl-CoA and Acetyl-CoA [66] (33),bt0388: from 2-Aminophenol derivative to *vic*-Dihydroxybenzenoid [19] (7, 26, 35),bt0391: From 3-Formiminopyruvate to 3-Formylpyruvate [22] by hydrolase (34), bt0021: From *cis*-2-Hydroxypenta-2,4-dienoate to 4-Hydroxy-2-oxovalerate [21] by hydratase (32), bt0022: Halomethyl derivative to 1-Methylalcohol derivative [23] (28), bt0082: from 2-Ketocarboxylate to Carboxylate [24] by oxidase (29).The results indicate neutral (yellow color) and likely transformations (green color), with no-toxic resultant products.




Figure 1. Biodegradation pathways of Isoniazid

Conclusion

Toxicological and biodegradability of the drug Isoniazid were evaluated by in silico predictions. Results showed that the drug able to degrade by microorganisms with different biotransformation reactions. The obtained results indicate the possibility to used drug with high precautions from environment and human health.

Acknowledgments

This work was supported by the Algerian Ministry of Higher Education and Scientific Research (MESRS) and the Thematic Agency for the Research in Health Sciences (ATRSS).

References

1. Pinedo-Rivilla, C., Aleu, J. and Collado, I.G. 2009. Pollutants biodegradation by fungi. *Current Organic Chemistry*, 13(12): 1194-1214.
2. Vorberg, S. and Tetko, I.V. 2014. Modeling the biodegradability of chemical compounds using the online CHEmical modeling environment (OCHEM). *Molecular Informatics*, 33(1): 73-85.
3. Vlădoiu, D.L., Filimon, M.N., Ostafe, V. and Isvoran, A. 2015. Effects of herbicides and fungicides on the soil chitinolytic activity. A molecular docking approach. *Ecological Chemistry and Engineering S*, 22(3): 439-450.

4. Książopolska, A. and Joniec, J. 2014. Activity of Microorganisms Participating in Organic Matter Transformation in Haplic Luvisol Soil. *Ecological Chemistry and Engineering S*, 21(3): 499-514.
5. Nishino, S.F. and Spain, J.C. 1995. Oxidative pathway for the biodegradation of nitrobenzene by *Comamonas sp.* strain JS765. *Applied and Environmental Microbiology*, 61(6): 2308-2313.
6. Gupta, R.C. and Shukla, O.P. 1978. Metabolism of isoniazid and related compounds by microorganisms: isolation, characterization and growth of isoniazid-degrading organism. *Indian Journal of Experimental Biology*, 16(10): 1047-1051.
7. Chauhan, A. and Jain, R.K. 2000. Degradation of o-nitrobenzoate via anthranilic acid (o-aminobenzoate) by *Arthrobacter protophormiae*: a plasmid-encoded new pathway. *Biochemical and Biophysical Research Communications*, 267(1): 236-244.
8. Noguera, D.R. and Freedman, D.L. 1996. Reduction and Acetylation of 2, 4-Dinitrotoluene by a *Pseudomonas aeruginosa* Strain. *Applied and Environmental Microbiology*, 62(7): 2257-2263.
9. Nishino, S.F. and Spain, J.C. 1993. Degradation of nitrobenzene by a *Pseudomonas pseudoalcaligenes*. *Applied and Environmental Microbiology*, 59(8): 2520-2525.
10. Preuss, A., Fimpel, J. and Diekert, G. 1993. Anaerobic transformation of 2, 4, 6-trinitrotoluene (TNT). *Archives of Microbiology*, 159(4): 345-353.
11. Gupta, R.C. and Shukla, O.P. 1979. Microbial transformation of isonicotinic acid hydrazide and isonicotinic acid by *Sarcina sp.* *Journal of Biosciences*, 1(2): 223-234.
12. Jang, J.Y., Kim, D., Bae, H.W., Choi, K.Y., Chae, J.C., Zylstra, G.J., Kim, Y.M. and Kim, E. 2005. Isolation and characterization of a *Rhodococcus species* strain able to grow on ortho-and para-xylene. *The Journal of Microbiology*, 43(4): 325-330.
13. Liu, Z., Yang, H., Huang, Z., Zhou, P. and Liu, S.J. 2002. Degradation of aniline by newly isolated, extremely aniline-tolerant *Delftia sp.* AN3. *Applied Microbiology and Biotechnology*, 58(5): 679-682.
14. Schenzle, A., Lenke, H., Fischer, P., Williams, P.A. and Knackmuss, H. 1997. Catabolism of 3-Nitrophenol by *Ralstonia eutropha* JMP 134. *Applied and Environmental Microbiology*, 63(4): 1421-1427.
15. Broderick, J.B. and O'Halloran, T.V. 1991. Overproduction, purification, and characterization of chlorocatechol dioxygenase, a non-heme iron dioxygenase with broad substrate tolerance. *Biochemistry*, 30(29): 7349-7358.
16. Toyama, T., Momotani, N., Ogata, Y., Miyamori, Y., Inoue, D., Sei, K., Mori, K., Kikuchi, S. and Ike, M. 2010. Isolation and characterization of 4-tert-butylphenol-utilizing *Sphingobium fuliginis* strains from *Phragmites australis* rhizosphere sediment. *Applied and Environmental Microbiology*, 76(20): 6733-6740.

17. Nelson, M.J. and Snell, E.E. 1986. Enzymes of vitamin B6 degradation. Purification and properties of 5-pyridoxic-acid oxygenase from *Arthrobacter sp.* Journal of Biological Chemistry, 261(32): 15115-15120.
18. Fuchs, A. and De Vries, F.W. 1978. Bacterial breakdown of benomyl. II. Mixed cultures. Antonie van Leeuwenhoek, 44(3-4): 293-309.
19. Rather, L.J., Knapp, B., Haehnel, W. and Fuchs, G. 2010. Coenzyme A dependent aerobic metabolism of benzoate via epoxide formation. Journal of Biological Chemistry, 285(27): 20615-2024.
20. Zhao, J.S., Singh, A., Huang, X.D. and Ward, O.P. 2000. Biotransformation of hydroxylaminobenzene and aminophenol by *Pseudomonas putida* 2NP8 cells grown in the presence of 3-nitrophenol. Applied and Environmental Microbiology, 66(6): 2336-2342.
21. Watson, G.K., Houghton, C. and Cain, R.B. 1974. Microbial metabolism of the pyridine ring. The metabolism of pyridine-3, 4-diol (3, 4-dihydroxypyridine) by *Agrobacterium sp.* Biochemical Journal, 140(2): 277-292.
22. Lau, P.C., Bergeron, H., Labbé, D., Wang, Y., Brousseau, R. and Gibson, D.T. 1994. Sequence and expression of the todGIH genes involved in the last three steps of toluene degradation by *Pseudomonas putida* F1. Gene, 146(1): 7-13.
23. Sette, L.D., Da Costa, L.A.M.A., Marsaioli, A.J. and Manfio, G.P. 2004. Biodegradation of alachlor by soil streptomycetes. Applied Microbiology and Biotechnology, 64(5): 712-717.
24. Blakley, E.R. 1977. The catabolism of L-tyrosine by an *Arthrobacter sp.* Canadian Journal of Microbiology, 23(9): 1128-1139.